



## Export Products

Product	Active ingredient	Formulation	Dosages
Fluocinoclear 750 Top Gel	Fluocinonide 0,075%	topic gel	30 g tube
Fluocinoclear Gel	Fluocinonide 0,025%	topic gel	30 g tube
Neobetaclear Gel	Betamethasone 0,05% + Neomicyn Sulphate 0,1%	topic gel	30 g tube
Clobetaderm White cream	Clobetasol 17-propionato 0,050%	topic cream	30 g tube
Neoclobetaderm White Cream	Clobetasol 17-propionato 0,05% + Neomicyn Sulphate 0,250%	topic cream	30 g tube
Betaderm White Cream	Betamethasone Dipropionate 0,064%	topic cream	30 g tube
Naltrexone Cloridrato	Naltrexone Chloridrate	1000 tablets	25 mg
Naltrexone Cloridrato	Naltrexone Chloridrate	1000 tablets	50 mg
Neopirin 100 mg	Acetylsalicylic Acyd	1000 tablets	100 mg
Neopirin 300 mg	Acetylsalicylic Acyd	1000 tablets	300 mg
Neopirin 500 mg	Acetylsalicylic Acyd	1000 tablets	500 mg
Rifampicina cps 300mg	Riphampicin	100 capsule	300 mg
Nitroxolina 50 mg	Nitroxoline	1000 tablets	50 mg
Nitroxolina 100 mg	Nitroxoline	1000 tablets	100 mg
Mebendazolo 100 mg	Mebendazole	1000 tablets	100 mg
Aminofillina 200 mg	Aminophylline	1000 tablets	200 mg
Metronidazolo 250 mg	Metronidazole	1000 tablets	250 mg
Potassio Permanganato 250 mg	Permanganate Potassium	1000 tablets	250 mg
Clorichina Difosfato 250 mg	Chloroquine Phosphate	1000 tablets	250 mg
Eritromicina 500 mg	Erythromicine	1000 tablets	500 mg
Lachitrim tablets	Trimethoprim 80 mg + Sulfametoxazolo 400 mg	1000 tablets	-
Lachitrim syrup	Trimethoprim 80 mg/100 ml + Sulfametoxazolo 400 mg/100 ml	25 bottles	-
Vitamine C	Ascorbic Acid	1000 tablets	250 mg
Vitamine C	Ascorbic Acid	1000 tablets	500 mg



## **Fluocinoclear 750 Top Gel Plus**

contains Fluocinonide 0,075% (75 mg for 100 gr).

### **ACTION AND USES**

The anti-inflammatory action of the active principle is increased in Fluocinoclear® 750 top gel plus, thanks to its higher dosage, that has been particularly studied in order to obtain the best and swiftest results without consequences or side-effects. Moreover, the particular compound of the gel allows an effective softening and smoothing action.

The use of Fluocinoclear® 750 top gel plus is therefore recommended in all inflammatory skin conditions, including eczema, dermatitis, psoriasis, and irritations in general. Smear small quantities on the affected area and massage till complete absorption is obtained.

### **DOSAGE**

Tube of 30 g.

One or two applications daily, according to the Physician's prescription.





## **Fluocinoclear Gel**

contains Fluocinonide 0,025% (25 mg for 100 gr).

### **ACTION AND USES**

Fluocinoclear® gel contains Fluocinonide, a very active topical corticosteroid, which is indicated for the treatment of a wide range of skin disorders as Dermatitis, Eczemas, Allergic reactions, Local inflammations, Prurigo, Psoriasis, etc.

Due to its particular composition Fluocinoclear® gel has emollient and lenitive effects and can be used on a wide area because it is well absorbed.

Massage a small quantity of Fluocinoclear® gel onto the skin until the complete absorption of the product.


### **DOSAGE**

Tube of 30 g.

Apply the gel one or two times daily, according to medical prescription.

**LACHIFARMA**





## **Neobetaclear Gel Plus**

contains Betamethasone 17,21 Dipropionate 64 mg (as Betamethasone 50 mg) Neomycin Sulphate 100mg 100gr.

### **ACTION AND USES**

Betamethasone is a powerful synthesis steroid and its ester, betamethasone dipropionate, is particularly suitable for dermatological use. Betamethasone exhibits a swift and strong local anti-inflammatory activity and is indicated in the treatment of all corticosteroid-sensitive dermatosis, while the addition of Neomycin Sulphate provides a broad spectrum antibacterial activity. Therefore, the use is recommended for the treatment of all dermatitis sensitive to topical steroids, exhibiting bacterial infections sensitive to neomycin, as well as of bacterial dermatitis showing evident inflammatory reactions. Massage the area to be treated till complete absorption is obtained.

**LACHIFARMA**





## **Clobetaderm White Cream**

contains Clobetasol Propionate 0,05%.

### **ACTION AND USES**

Clobetaderm® white cream is a very active topical medicament and is particularly effective as anti-inflammatory, antiallergic and antipruritic. Therefore, the use of Clobetaderm® white cream is recommended for the treatment of psoriasis, recalcitrant eczemas, lichen planus, discoid lupus erythematosus and other conditions which do not respond satisfactorily to less active steroids.

### **DOSAGE**

Tube of 30 g.

Apply sparingly the affected area once or twice daily until improvement occurs. As with other highly-active steroid preparations, therapy should be discontinued when control is achieved.

**LACHIFARMA**





## **Neoclobetaderm White Cream**

contains Neomycin Sulphate 0,25% and Clobetasol Propionate 0,05%.

### **ACTION AND USES**

Neoclobetaderm ® white cream is a very active topical medicament and is particularly effective against resistant dermatoses, especially if accompanied by skin infections, except if caused by viruses, fungi or bacteria. Therefore, the use of Neoclobetaderm ® white cream is recommended for the treatment of psoriasis, eczemas, lichen planus, discoid lupus erythematosus, wherever such conditions do not respond to less active steroids satisfactorily.

### **DOSAGE**

Tube of 30 g.

Apply to the affected area once or twice daily, as suggested by the Physician.

**LACHIFARMA**





## **Betaderm White Cream**

contains Betamethasone 0,05% (Dipropionate)

### **ACTION AND USES**

Betaderm® white cream contains Betamethasone 17,21 Dipropionate, an effective topical corticosteroid with a special anti-inflammatory and anti-allergic action. Betaderm ® white cream is suitable for the treatment of a wide range of skin disorders: infective eczemas, Psoriasis, Lichen Ruber Planus, Prurigo, Lood inflammations etc.. Thanks to its particular composition, Betaderm white cream is well absorbed and may be utilized on a wide area.

### **DOSAGE**

Tube of 30 g.

Apply a small quantity of Betaderm ® white cream and massage it until the product is completely absorbed. Adopt one or two applications daily, subject to medician's, prescription.

**LACHIFARMA**

